

$$\int_0^{\infty} e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$$

James T. Smith
San Francisco State University

Define

$$g(x) = \int_0^x e^{-t^2} dt$$

for all real x . If $1 \leq t \leq x$, then $t \leq t^2$, so $e^{-t^2} \leq e^{-t}$ and

$$0 \leq \int_1^x e^{-t^2} dt \leq \int_1^x e^{-t} dt = -e^{-t} \Big|_{t=1}^x = -e^{-x} + e^{-1}.$$

Since $e^{-x} \rightarrow 0$ as $x \rightarrow \infty$,

$$J = \int_0^{\infty} e^{-t^2} dt = \lim_{x \rightarrow \infty} g(x) = g(1) + \lim_{x \rightarrow \infty} \int_1^x e^{-t^2} dt$$

exists. By the fundamental theorem of calculus and the chain rule,

$$\begin{aligned} g'(t) &= e^{-t^2} \\ \frac{d}{dx} g(\sqrt{x}) &= 2g(\sqrt{x})g'(\sqrt{x}) \frac{1}{2\sqrt{x}} = \frac{e^{-x}}{\sqrt{x}} g(\sqrt{x}) \\ &= \frac{e^{-x}}{\sqrt{x}} \int_0^{\sqrt{x}} e^{-t^2} dt = e^{-x} \int_0^1 e^{-xu^2} du \end{aligned}$$

provided $x > 0$, where $t = u\sqrt{x}$.

Now consider the function

$$f(x) = \int_0^1 \frac{e^{-x(1+u^2)}}{1+u^2} du$$

By a standard calculus theorem about single integrals of multivariate functions,

$$f'(x) = \int_0^1 \frac{-(1+u^2)e^{-x(1+u^2)}}{1+u^2} du = -\int_0^1 e^{-x-xu^2} du = -e^{-x} \int_0^1 e^{-xu^2} du$$

for all x . Moreover,

$$\int_0^{\infty} e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$$

$$f(0) = \int_0^1 \frac{1}{1+u^2} du = \arctan 1 - \arctan 0 = \frac{\pi}{4},$$

f is continuous, and

$$\lim_{x \rightarrow \infty} f(x) = \int_0^1 \lim_{x \rightarrow \infty} \frac{e^{-x(1+t^2)}}{1+t^2} dt = \int_0^1 0 dt = 0.$$

Since

$$f'(x) = -\frac{d}{dx} g(\sqrt{x})^2,$$

$f(x) + g(\sqrt{x})^2$ is constant for all $x > 0$. Since g is continuous too, that constant is

$$\lim_{x \rightarrow 0} \left(f(x) + g(\sqrt{x})^2 \right) = f(0) + g(0)^2 = \frac{\pi}{4}.$$

It follows that

$$\frac{\pi}{4} = \lim_{x \rightarrow \infty} \left(f(x) + g(\sqrt{x})^2 \right) = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(\sqrt{x})^2 = 0 + J^2.$$

Thus

$$J = \frac{\sqrt{\pi}}{2}.$$

References

Robert WEINSTOCK, “Elementary evaluations of $\int_0^{\infty} e^{-x^2} dx$, $\int_0^{\infty} \cos x^2 dx$, and $\int_0^{\infty} \sin x^2 dx$,” *American mathematical monthly*, XCVII (1990), 39-42. See also the literature cited there. [Click here](#) to see the article.

Constantine GEORGAKIS, “A note on the Gaussian integral,” *Mathematics magazine*, LXVII (1994), 47. An alternate approach, easier to motivate but harder to justify.